

Elyxyb[®]
(celecoxib)
Oral Solution

**THINK
FAST**

**EXTINGUISH YOUR
MIGRAINE PAIN**

FAST

WITH Elyxyb[®]
(celecoxib)
Oral Solution

If you're not getting the relief you
need from your current medication

**LOOK
INSIDE**

APPROVED USE

ELYXYB[®] (celecoxib) oral solution is:

A prescription nonsteroidal anti-inflammatory drug (NSAID) used for the acute treatment of migraine with or without aura in adults.

- ELYXYB is not used as a preventive treatment of migraine
- It is not known if ELYXYB is safe and effective in children

IMPORTANT SAFETY INFORMATION about ELYXYB

WARNING: RISK OF SERIOUS CARDIOVASCULAR and GASTROINTESTINAL EVENTS

Cardiovascular Thrombotic Events

- Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in the treatment and may increase with duration of use
- ELYXYB is contraindicated in the setting of coronary artery bypass graft (CABG) surgery

Gastrointestinal Bleeding, Ulceration, and Perforation

- NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious (GI) events

Please see Important Safety Information on the next page and full [Prescribing Information](#), including Boxed Warning.

Do you feel like you're waiting for migraine relief?

Migraine pain can feel like a fire going on in your head, and for some, the fire is still there even after taking migraine medication. *If you feel like this, your current medication may not be giving you the relief you need.*

If any of these are true for you, you may need something different:



Do you still have migraine pain 2 hours after you take your medication?



Does your migraine come back within 24 hours?



Have you ever stopped taking your migraine medication or thought twice about taking it because you didn't want the drowsiness, nausea, or other side effects?



Talk to your doctor about ELYXYB today

IMPORTANT SAFETY INFORMATION about ELYXYB (cont'd)

Do not take ELYXYB:

- if you are allergic to celecoxib or any of the ingredients in ELYXYB
- if you are allergic to sulfonamides
- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs
- right before or after heart bypass surgery

Please see Important Safety Information continued on the next page and full [Prescribing Information](#), including [Boxed Warning](#).

Extinguish your migraine pain *FAST* with ELYXYB

In 2 clinical studies, ELYXYB

DELIVERED PAIN RELIEF IN AS SOON AS 15 MINUTES



1 HOUR

MAJORITY (over 5 out of 10) of people felt **PAIN RELIEF***

2 HOURS

MANY (over 1 out of 3) people were **100% PAIN FREE†**

24 HOURS

NEARLY EVERYONE (over 9 out of 10) continued to be **100% PAIN FREE‡** through 24 hours‡

*vs 42.5% of people on placebo.

†vs 24% of people on placebo.

‡vs 88% of people on placebo.

Relieved other migraine symptoms:

Nausea

Sensitivity to light

Sensitivity to sound

No one in clinical trials stopped taking ELYXYB because of drowsiness, nausea, or other treatment-related side effects.

Drowsiness was rare and no serious side effects occurred during clinical studies. The most common side effect people reported with ELYXYB was food tasting differently.

IMPORTANT SAFETY INFORMATION about ELYXYB (cont'd)

ELYXYB should only be used:

- exactly as prescribed
- for the shortest time needed

Please see Important Safety Information continued on the next page and full **Prescribing Information**, including **Boxed Warning**.

Elyxyb® | **THINK**
(celecoxib) | **FAST**
Oral Solution

You could save on your
ELYXYB prescription*

- 1 Ask your doctor to prescribe ELYXYB
- 2 Present this copay savings card to your pharmacist
- 3 Pay as little as \$0 on each prescription and \$25 for refills*
- 4 Ask your doctor for FREE SAMPLES of ELYXYB

Patient Access Support & Savings Program

Elyxyb® (celecoxib)
Oral Solution
Passport

BIN# 600426
PCN# 54
GRP# EC45401001
ID# 39854504969

Pay as little as **\$0**
on each prescription
and **\$0** for refills*

*Commercial insurance patients only.
Terms and conditions apply.

THIS IS AN ACTIVE CARD FOR YOUR PHARMACIST

*For commercial patients only; terms and conditions apply.

Please see Important Safety Information continued on the next page and full **Prescribing Information**, including Boxed Warning.

See how easy fast relief can be

Just 1 dose stops migraine

– Don't take more than 1 dose per day

Ready-to-drink bottles: no mixing, shaking, or measuring needed

– Less than 1 teaspoon dose

Take with or without food

– After taking, you may drink up to 8 ounces (1 cup) of water

Convenient, on-the-go treatment

– A small bottle that fits in pockets and purses



Not actual size.

Elyxyb®
(celecoxib)
Oral Solution

**THINK
FAST**

When migraine hits, every minute matters. Ask your doctor about ELYXYB

IMPORTANT SAFETY INFORMATION about ELYXYB (cont'd)

ELYXYB contains celecoxib, a NSAID. NSAIDs, including ELYXYB, can cause serious side effects, including:

- **Increased risk of a heart attack or stroke that can lead to death. This risk may happen early in treatment and may increase:**

- with increasing doses of NSAIDs
- with longer use of NSAIDs

- **Do not take ELYXYB right before or after a heart surgery called a “coronary artery bypass graft (CABG).”**

Avoid taking NSAIDs, including ELYXYB, after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.

- **Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach, and intestines:**

- anytime during use
- without warning symptoms
- that may cause death

Don't start a new medication before talking to your healthcare provider and telling them about all of the medications you take.

Before taking ELYXYB, tell your healthcare provider about all of your medical conditions, including if you:

- have liver or kidney problems
- have a history of stomach ulcer or bleeding in your stomach or intestines
- have heart disease or risk factors that increase your chance of getting heart disease
- have high blood pressure

- have asthma
- are pregnant or plan to become pregnant. Taking NSAIDs, including ELYXYB, at about 20 weeks of pregnancy or later may harm your unborn baby. If you need to take NSAIDs for more than 2 days when you are between 20 and 30 weeks of pregnancy, your healthcare provider may need to monitor the amount of fluid in your womb around your baby. **You should not take NSAIDs after about 30 weeks of pregnancy**
- are breastfeeding or plan to breastfeed. ELYXYB may pass into your breast milk. Talk with your healthcare provider about the best way to feed your baby if you take ELYXYB

When taking ELYXYB

See the detailed “Instructions for Use” on how to take ELYXYB solution.

- Take ELYXYB exactly as your healthcare provider tells you to take it
- Take ELYXYB by mouth with or without food
- Do not take more than one dose in a 24-hour period
- Use ELYXYB for the fewest number of days a month, as needed

The possible side effects of ELYXYB are:

liver problems including liver failure, new or worse high blood pressure, heart failure, kidney problems including kidney failure, life-threatening allergic reactions, asthma attacks in people who have asthma, life-threatening skin reactions, medication overuse headaches, low red blood cells (anemia). **Other side effects of NSAIDs include:** stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

Stop taking ELYXYB and call your healthcare provider right away if you get any of the following symptoms:

- nausea
- more tired or weaker than usual
- diarrhea
- itching
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms, legs, hands and feet

Get emergency help right away if you have:

Shortness of breath or trouble breathing, chest pain, weakness in one part or side of your body, slurred speech, or swelling of the face or throat.

If you take too much ELYXYB, call your healthcare provider or get medical help right away.

These are not all the side effects associated with ELYXYB. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

For more information, go to dailymed.nlm.nih.gov.

Please see full [Prescribing Information](#), including [Boxed Warning](#).



ELYXYB® is a registered trademark of SCILEX Pharmaceuticals Inc. All other trademarks are the property of their respective owners. © 2024 SCILEX Holding Company. All rights reserved. ELY-00174. 6/24

